

of diabetes. These effects may be mediated by the direct effects of insulin on the myocardium and coronary vasculature or indirectly through systemic reductions in blood glucose. Anti-angiogenic proteins, angiostatin and endostatin, are downregulated with insulin treatment and appear to play an important role in the diabetic angiogenic response. Insulin therapy appears to be a promising strategy to improve the results of growth factor or cell-based angiogenic therapies in diabetic patients.

## References

1. Simons M, Bonow RO, Chronos NA, Cohen DJ, Giordano FJ, Hammond HK, et al. Clinical trials in coronary angiogenesis: issues, problems, consensus—an expert panel summary. *Circulation*. 2000;102:E73-86.
2. Boodhwani M, Sodha NR, Laham RJ, Sellke FW. The future of therapeutic myocardial angiogenesis. *Shock*. 2006;26:332-41.
3. Boodhwani M, Nakai Y, Mieno S, Voisine P, Bianchi C, Araujo EG, et al. Hypercholesterolemia impairs the myocardial angiogenic response in a swine model of chronic ischemia: role of endostatin and oxidative stress. *Ann Thorac Surg*. 2006;81:634-41.
4. Ruel M, Wu GF, Khan TA, Voisine P, Bianchi C, Li J, Laham RJ, et al. Inhibition of the cardiac angiogenic response to surgical FGF-2 therapy in a swine endothelial dysfunction model. *Circulation*. 2003;108(Suppl 1):II335-40.
5. Voisine P, Bianchi C, Ruel M, Malik T, Rosinberg A, Feng J, et al. Inhibition of the cardiac angiogenic response to exogenous vascular endothelial growth factor. *Surgery*. 2004;136:407-15.
6. Mieno S, Boodhwani M, Clements RT, Ramlawi B, Sodha NR, Li J, et al. Aging is associated with an impaired coronary microvascular response to vascular endothelial growth factor in patients. *J Thorac Cardiovasc Surg*. 2006;132:1348-55.
7. Zimmet P, Alberti KG, Shaw J. Global and societal implications of the diabetes epidemic. *Nature*. 2001;414:782-7.
8. Abaci A, Oguzhan A, Kahraman S, Eryol NK, Unal S, Arinc H, et al. Effect of diabetes mellitus on formation of coronary collateral vessels. *Circulation*. 1999;99:2239-42.
9. Yarom R, Zirkun H, Stammler G, Rose AG. Human coronary microvessels in diabetes and ischaemia. Morphometric study of autopsy material. *J Pathol*. 1992;166:265-70.
10. Boodhwani M, Sodha NR, Mieno S, Xu S, Feng J, Ramlawi B, et al. Functional, cellular, and molecular characterization of the angiogenic response to chronic myocardial ischemia in diabetes. *Circulation*. In press.
11. Tofukuji M, Metais C, Li J, Hariawala MD, Franklin A, Vassileva C, et al. Effects of ischemic preconditioning on myocardial perfusion, function, and microvascular regulation. *Circulation*. 1998;98(19 Suppl):II197-204; discussion II204-5.
12. The effect of intensive treatment of diabetes on the development and progression of long-term complications in insulin-dependent diabetes mellitus. The Diabetes Control and Complications Trial Research Group. *N Engl J Med*. 1993;329:977-86.
13. Standards of medical care in diabetes. *Diabetes Care*. 2005;28:S4-36.
14. Brownsey RW, Boone AN, Allard MF. Actions of insulin on the mammalian heart: metabolism, pathology and biochemical mechanisms. *Cardiovasc Res*. 1997;34:3-24.
15. White MF. Insulin signaling in health and disease. *Science*. 2003;302:1710-1.
16. Zhang L, Zalewski A, Liu Y, Mazurek T, Cowan S, Martin JL, et al. Diabetes-induced oxidative stress and low-grade inflammation in porcine coronary arteries. *Circulation*. 2003;108:472-8.
17. Wautier JL, Schmidt AM. Protein glycation: a firm link to endothelial cell dysfunction. *Circ Res*. 2004;95:233-8.
18. Sharma MR, Tuszynski GP, Sharma MC. Angiostatin-induced inhibition of endothelial cell proliferation/apoptosis is associated with the down-regulation of cell cycle regulatory protein cdk5. *J Cell Biochem*. 2004;91:398-409.
19. O'Reilly MS, Boehm T, Shing Y, Fukai N, Vasios G, Lane WS, et al. Endostatin: an endogenous inhibitor of angiogenesis and tumor growth. *Cell*. 1997;88:277-85.
20. Weihrauch D, Lohr NL, Mraovic B, Ludwig LM, Chilian WM, Pagel PS, et al. Chronic hyperglycemia attenuates coronary collateral development and impairs proliferative properties of myocardial interstitial fluid by production of angiostatin. *Circulation*. 2004;109:2343-8.
21. Chung AW, Hsiang YN, Matzke LA, McManus BM, van Breemen C, Okon EB. Reduced expression of vascular endothelial growth factor paralleled with the increased angiostatin expression resulting from the upregulated activities of matrix metalloproteinase-2 and -9 in human type 2 diabetic arterial vasculature. *Circ Res*. 2006;99:140-8.
22. Matsunaga T, Chilian WM, March K. Angiostatin is negatively associated with coronary collateral growth in patients with coronary artery disease. *Am J Physiol Heart Circ Physiol*. 2005;288:H2042-6.
23. Guo M, Wu MH, Korompai F, Yuan SY. Upregulation of PKC genes and isozymes in cardiovascular tissues during early stages of experimental diabetes. *Physiol Genomics*. 2003;12:139-46.

## Discussion

**Dr Mark J. Krasna (Towson, Md).** I was quite impressed with the technical feat of this study. Can you elaborate on the three procedures that you performed on each of these animals and was there any mortality among the three groups?

**Dr Boodhwani.** In the past we have done more than 400 ameroid constrictor experiments in our laboratory. We found that in a potent model of diet induced hypercholesterolemia, there was about a 10% to 20% mortality rate with the animals, but over time the techniques have been improved and we have a much better sense of what the model entails now. Fortunately, in this study, we did not lose any animals.

**Dr Krasna.** Along the same lines, when you are doing three procedures on each animal, was there a significant infection rate from the first to the third, or was that also low?

**Dr. Boodhwani.** In the normal, nondiabetic animals (group ND) there was virtually no infection. In the diabetic animals we found that there was some impairment of wound healing and breakdown of the incision. However, to emphasize, all animals were treated with postoperative antibiotics for 5 days.

**Dr Paul W. M. Fedak (Calgary, Alberta, Canada).** This is a great study. I think this model is going to be very important for future studies in the area. My question to you is, I did not see anything about regional or global left ventricular function. Was that addressed in this study or will it be addressed in any future studies? If not, what would you hypothesize would be the effects of insulin treatment?

**Dr Boodhwani.** In the past we have demonstrated that circumflex ameroid constrictor placement leads to a reduction in perfusion as well as regional wall motion abnormalities. Regional wall motion abnormalities were not specifically studied in this model, although we did assess global left ventricular function. We expect, though, that improvement in perfusion would likely result in improvement in regional wall motion as well.